



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).


57. Proposed by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics in Russell College, Lebanon, Va.

A chord is drawn through two points taken at random in the surface of a circle. If a second chord be drawn through two other points taken at random in the surface, find the chance that the quadrilateral formed by joining the extremities of the two chords will contain the center of the circle.

I. Solution by the PROPOSER.

Let  $M, N$  be the first two random points;  $AB$  the chord through them;  $P, Q$  the second two random points;  $CD$  the chord through them; and  $O$  the center of the circle. Draw  $OH, OK$  perpendicular to  $AB, CD$ .

Let  $AO=r, AM=w, MN=x, CP=y, PQ=z, \angle AOH=\theta, \angle COK=\varphi, \angle AOC=\psi$ , and  $\mu$ =the angle  $AB$  makes with some fixed line.



Then  $AH=r\sin\theta, CK=r\sin\varphi$ ; an element of the circle at  $M$  is  $r\sin\theta d\theta dw$ ; at  $N, d\mu x dx$ ; at  $P, r\sin\varphi d\varphi dy$ ; at  $Q, d\psi z dz$ . The limits of  $\theta$  are 0 and  $\frac{1}{2}\pi$ ; of  $\varphi, \theta$  and  $\theta$ , and doubled; of  $\psi, \pi-2\theta$  and  $\pi$ , and doubled; of  $\mu, 0$  and  $2\pi$ ; of  $w, 0$  and  $2r\sin\theta=s$ ; of  $x, 0$  and  $w$ , and doubled; of  $y, 0$  and  $2r\sin\varphi=v$ ; and of  $z, 0$  and  $y$ , and doubled. Hence, since the whole number of ways the four points can be taken is  $\pi^4 r^8$ , the required chance is

$$\begin{aligned}
 p &= \frac{16}{\pi^4 r^8} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\pi-2\theta}^\pi \int_0^{2\pi} \int_0^s \int_0^w \int_0^v \int_0^y r\sin\theta d\theta r\sin\varphi d\varphi d\psi d\mu dw dx dy dz \\
 &= \frac{8}{\pi^4 r^6} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\pi-2\theta}^\pi \int_0^{2\pi} \int_0^s \int_0^w \int_0^v \sin\theta \sin\varphi d\theta d\varphi d\psi d\mu dw dx dy^2 dz \\
 &= \frac{64}{3\pi^4 r^3} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\pi-2\theta}^\pi \int_0^{2\pi} \int_0^s \int_0^w \sin\theta \sin^4 \varphi d\theta d\varphi d\psi d\mu dw dx \\
 &= \frac{32}{3\pi^4 r^3} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\pi-2\theta}^\pi \int_0^{2\pi} \int_0^s \sin\theta \sin^4 \varphi d\theta d\varphi d\psi d\mu w^2 dw \\
 &= \frac{256}{9\pi^4} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\pi-2\theta}^\pi \int_0^{2\pi} \sin^4 \theta \sin^4 \varphi d\theta d\varphi d\psi d\mu \\
 &= \frac{512}{9\pi^3} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_{\pi-2\theta}^\pi \sin^4 \theta \sin^4 \varphi d\theta d\varphi d\psi = \frac{1024}{9\pi^3} \int_0^{\frac{1}{2}\pi} \int_0^\theta \theta \sin^4 \theta \sin^4 \varphi d\theta d\varphi \\
 &= \frac{128}{9\pi^3} \int_0^{\frac{1}{2}\pi} (3\theta^2 - 3\theta \sin\theta \cos\theta - 2\theta \sin^3 \theta \cos\theta) \sin^4 \theta d\theta = \frac{2}{3} + \frac{139}{72\pi^2}.
 \end{aligned}$$

II. Solution by HENRY HEATON, M. Sc., Atlantic, Iowa.

Let  $A$  and  $D$  represent the positions of the first and third points, and sup-

pose the second and fourth to be taken somewhere upon the lines  $BC$  and  $EF$ , respectively.

Put  $OC=a$ ,  $OA=x$ ,  $OD=y$ ,  $\angle OAC=\theta$ ,  $\angle IOC=\phi$ ,  $\angle ODK=\psi$ , and  $\angle KOF=\omega$ .

Then  $a\cos\phi=x\sin\theta$ , and  $a\cos\omega=y\sin\psi$ .

If either  $E$  or  $F$  is upon the arc  $HG$ , the quadrilateral formed by joining the extremities of the chords will contain the center of the circle.

Since the arc  $HG$ =the arc  $CB$ , the probability of this is  $(\phi+\omega)/\pi$ .

If  $x>0$  and  $<a$  the chance that the first point is taken between  $x$  and  $x+dx$  is  $2\pi x dx/a^2\pi=2xdx/a^2$ .

If  $y>0$  and  $<a$  the chance that the third point is taken between the distances  $y$  and  $y+dy$  is  $2ydy/a^2$ .

If  $\theta>0$  and  $<\frac{1}{2}\pi$  the chance that the second point is taken between the line  $BC$  and a second line making at  $A$  the angle  $d\theta$  is  $\frac{1}{2}(AC^2+AB^2)d\theta/\frac{1}{2}a^2\pi=2[a^2+x^2(1-2\sin^2\theta)]d\theta/a^2\pi=2(1-2\cos^2\phi+\cos^2\phi\operatorname{cosec}^2\theta)d\theta/\pi$ .

If  $\psi>0$  and  $<\frac{1}{2}\pi$  the chance that the fourth point is taken between the line  $EF$  and a second line making at  $D$  the angle  $d\psi$  is  $2(1-2\cos^2\omega+\cos^2\omega\operatorname{cosec}^2\psi)d\psi/\pi$ .

If we suppose  $\theta$  constant while  $x$  and  $\phi$  vary,  $xdx=-a^2\sin\phi\cos\phi\operatorname{cosec}^2\theta d\phi$ . When  $x=0$ ,  $\phi=\frac{1}{2}\pi$ , and when  $x=a$ ,  $\phi=\frac{1}{2}\pi-\theta$ . Hence the limits of integration for  $\phi$  are  $\frac{1}{2}\pi-\theta$  and  $\frac{1}{2}\pi$ . If we integrate first with respect to  $\theta$ , the limits for  $\theta$  are  $\frac{1}{2}\pi-\phi$  and  $\frac{1}{2}\pi$ , while for  $\phi$  they are 0 and  $\frac{1}{2}\pi$ .

In like manner we may substitute  $-a^2\sin\omega\cos\omega\operatorname{cosec}^2\psi d\omega$  for  $ydy$  and integrate between limits  $\frac{1}{2}\pi-\omega$  and  $\frac{1}{2}\pi$  for  $\psi$ , and between 0 and  $\frac{1}{2}\pi$  for  $\omega$ .

Hence the required probability is

$$\begin{aligned}
 P &= \frac{16}{\pi^3} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_{\frac{1}{2}\pi-\phi}^{\frac{1}{2}\pi} \int_{\frac{1}{2}\pi-\omega}^{\frac{1}{2}\pi} (\phi+\omega)(1-2\cos^2\phi+\cos^2\phi\operatorname{cosec}^2\theta) \times \\
 &\quad (1-2\cos^2\omega+\cos^2\omega\operatorname{cosec}^2\psi) \sin\phi\cos\phi\operatorname{cosec}^2\theta \sin\omega\cos\omega\operatorname{cosec}^2\psi d\phi d\omega d\theta d\psi \\
 &= \frac{256}{9\pi^3} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} (\phi+\omega) \sin^4\phi \sin^4\omega d\phi d\omega \\
 &= \frac{4}{9\pi^3} \int_0^{\frac{1}{2}\pi} (12\pi\phi+3\pi^2+16)\sin^4\phi d\phi = \frac{1}{2} + (8/3\pi^2) = .77019.
 \end{aligned}$$

## MISCELLANEOUS.

55. Proposed by J. M. COLAW, A. M., Monterey, Va.

Multiply 6 by 4. Is the problem legitimate when both symbols represent pure number?

[NOTE. "A measured or numbered quantity may be divided into a number of parts, or taken a number of times; but no number can be multiplied or divided into parts."—*McClellan and Dewey's Psychology of Number*. "The astounding thesis is maintained that number is not a magnitude, does not possess quantity at all, and that 'no number can be multiplied or divided into parts'."—*Lefevre's Number and Its Algebra*.]

